



SuperFix

Bactericide/Fungicide

Guaranteed Analysis

Active Ingredients:

Clove 12.5%, Cottonseed oil 11.5%, Rosemary 10.6%, Thyme 10.2%, Peppermint oil 9.5%, Cinnamon 8.2%, Garlic 8.2% (71.7%)

Other Ingredients:

White Mineral Oil, Palm Oil, Guar Gum, Lemon Juice Concentrate, Citrus pulp - (29.3%)

Total - 100%

CAUTION

PRECAUTIONARY STATEMENTS

May cause temporary eye and skin irritation. Do not get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses, and chemical resistant gloves. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If irritation persists, call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing, rinse skin immediately with plenty of soap and water for 15-20 minutes. If irritation persists, call a poison control center or doctor for further advice.

If swallowed: Call doctor or poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.

DIRECTIONS FOR USE

APPLICATION GUIDELINES. SuperFix does not normally require a pH change. It is recommended that a pH in the range of 7.0 or less be used. SuperFix is a multi-purpose broad spectrum bactericide/fungicide designed for cannabis crop. SuperFix can be used for soil applications to aid in reducing the damaging effects of soil borne bacterial and fungal diseases.

APPLICATION TIMING. Apply diluted spray when high humidity for the development of disease symptoms in crops or at first sign of disease and continue applications at 7 to 10 day intervals or until disease symptoms are gone.

MINIMUM TIME FOR APPLICATION TO HARVEST. Spraying or drip application of the product can occur up to the time of harvest.

APPLICATION METHOD AND EQUIPMENT. SuperFix can be sprayed on foliage, the surface of soil or injected into a drip irrigation system at the rates as shown. When sprayed on foliage allow 12 hours of a dry period so that irrigation or rain does not wash off application. For drip applications apply to soil that is with humidity level of 70% or more. After SuperFix has cleared lines follow with enough water to wash down to root level. Always wash into soil after application with plain water to make sure that SuperFix penetrates down to the depth of the root ball. Thorough coverage of foliage is required for optimal disease control. Use adequate sprayer pressure to achieve small droplets or fine mist for best coverage. Do not spray wet foliage.

STORAGE AND DISPOSAL. Store in original UV protected containers only. In case of spill on floor or paved surfaces, hose off with clean water. Store at temperatures between 41°F to 85°F (5°C to 30°C). Wastes resulting from using this product may be disposed of on site. Dispose of empty containers in a sanitary land fill or recycled if permitted.

CONDITIONS OF SALE AND WARRANTY

Mar Vista Resources LLC, (Seller) warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon Seller. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition, of sale by Seller and is accepted as such by the buyer. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of Seller's liability.

Crop	Use Rate per 100 gallons	Application Rates and Timing	Diseases and Bacteria
Cannabis Hemp	32 fl. Oz to 128 fl. Oz.	Use lower rate when pest and disease pressure is light and spray at 10-14 day intervals. During conditions of high pest or disease pressure increase rates or shorten intervals to 7-10 days.	Fusarium wilt, Fusarium stem canker, Fusarium root rot, Rhizopus, Powdery mildew, Phytoplasma bacteria, Pythium crown rot, Leaf blight, Alternaria, Phytophthora

Minimum Risk Pesticides Exempted under FIFRA Section 25(b)



- ☐ NET CONTENTS: 1 Qt (0.946 L) Jug
- ☐ NET CONTENTS: 1 Gal (3.69 L) Jug
- ☐ NET CONTENTS: 55 gal (208.19 L) Drum



Distributed By:

Mar Vista Resources, LLC at 745 North Ave • Corcoran, CA 93212 • Office: (559) 992-4535

Revision: 06-2020 GHS